



US 20140323763A1

(19) **United States**

(12) **Patent Application Publication**

Furkert et al.

(10) **Pub. No.: US 2014/0323763 A1**

(43) **Pub. Date: Oct. 30, 2014**

(54) **ANTIBACTERIAL AND ANTIFUNGAL
SUBSTANCES BIPHENYLYL COMPOUNDS**

Publication Classification

(75) Inventors: **Franz Furkert**, Preetz (DE); **Bernd
Clement**, Kiel (DE); **Britta Gerig**,
Munchen (DE); **Dieter Heber**, Molfsee
(DE)

(51) **Int. Cl.**
C07C 225/16 (2006.01)
C07C 211/63 (2006.01)
C07C 211/29 (2006.01)
(52) **U.S. Cl.**
CPC *C07C 225/16* (2013.01); *C07C 211/29*
(2013.01); *C07C 211/63* (2013.01)
USPC **564/289**; 564/337

(73) Assignee: **CHRISTIAN-ALBRECHTS-
UNIVERSITY OF KIEL**, Kiel (DE)

(21) Appl. No.: **13/995,261**

(57) **ABSTRACT**

(22) PCT Filed: **Dec. 21, 2011**

(86) PCT No.: **PCT/EP2011/073601**

§ 371 (c)(1),

(2), (4) Date: **Aug. 30, 2013**

The invention relates to the use of a substance of the general form (I) to produce an antibacterial and/or antifungal drug, wherein X is a methylene group or a carbonyl group; R₁, R₂, and R₃ are each selected from the group comprising hydrogen, an alkyl group having a chain length of 1-4 carbon atoms, an alkoxy group having a chain length of 1-3 carbon atoms, and a halogen; R₄ and R₅ are each selected from the group comprising hydrogen and an alkyl group having a chain length of 1-4 carbon atoms; and n=3 to 6.

(30) **Foreign Application Priority Data**

Dec. 21, 2010 (DE) 10 2010 055 322.0

ANTIBACTERIAL AND ANTIFUNGAL SUBSTANCES BIPHENYLYL COMPOUNDS

[0001] The invention relates to antibacterial and antimycotic active substances and their use for production of pharmaceutical compositions.


[0002] Despite all scientific progress bacterial infections are a constant threat to mankind.

[0003] Today's threat is primarily due to the increase of resistance to conventional antibiotics and the low number of new drugs [Gould, I. M., "Antibiotic resistance: the perfect storm" International Journal of Antimicrobial Agents, 2009, 34 (3), 2-5].

[0004] Methicillin-resistant germs represent a specific problem; if a *Staphylococcus aureus* is methicillin-resistant (so-called MRSA) often resistances against antibiotics of other groups (quinolones, tetracyclines, aminoglycosides, erythromycins, sulfonamides) are displayed.

[0005] Therefore, the problem underlying the present invention is to provide new compounds which are effective against various bacteria and fungi, especially against those which are already showing multidrug resistance and in this context are suitable as pharmaceutical agents.

[0006] It has been shown that basic biphenyls of the general formula



wherein X is a methylene group (CH₂) or a carbonyl group (C=O), R₁, R₂ and R₃ are hydrogen, an alkyl group having a

chain length of 1-4 carbon atoms, (e.g. methyl (CH₃)—, ethyl (CH₂CH₃)—, propyl (CH₂CH₂CH₃)—, isopropyl (CH(CH₃)CH₂)—, butyl (CH₂CH₂CH₂CH₃)—, isobutyl (CH(CH₃)CH₂CH₃)— or tert-butyl (C(CH₃)₃)— group), an alkoxy group having a chain length of 1-3 carbon atoms (e.g. a methoxy (OCH₃)—, ethoxy (OCH₂CH₃)—, propoxy (OCH₂CH₂CH₃)— or isopropoxy (OCH(CH₃)CH₂)— group) or a halogen (e.g. Fluorine (F), chlorine (Cl), bromine (Br) or iodine (I)),

R₄ and R₅ are hydrogen or an alkyl group having a chain length of 1-4 carbon atoms (e.g. methyl (CH₃)—, ethyl (CH₂CH₃)—, propyl (CH₂CH₂CH₃)—, isopropyl (CH(CH₃)CH₂)—, butyl (CH₂CH₂CH₂CH₃)—, isobutyl (CH(CH₃)CH₂CH₃)—, or tert-Butyl (C(CH₃)₃)— group) and

wherein n can be n=3-6, as well as salts, hydrochlorides and tautomers of those compounds have an antimicrobial and antifungal activity also against methicillin-resistant germs. Therefore, the bisphenyls as described above and illustrated in the following embodiments are suitable for inhibiting the growth of bacteria and/or fungi. Accordingly, the present invention relates to the use of the substances as antibacterial and/or antifungal drugs.

[0007] Table 1 shows the values of the minimum inhibitory concentration (MIC) of the substances of the present invention. For comparison, the substance GG20-3, which differs from the substances of the present invention by a shortened methyl group chain (n=2) and the known antibiotics vancomycin and tetracycline, were also measured.

Determination of the minimum inhibitory concentration (MIC)

[0008] The minimum inhibitory concentration is the lowest concentration of the tested substance which completely inhibits the growth of each test organism, i.e. wherein after incubation the turbidity is not measurable. The relevant determination is valid if the growth control (DMSO and medium) a distinct turbidity is shown.

Abbreviations

[0009] *Staphylococcus aureus* S. a., methicillin-resistant *Staphylococcus aureus* MRSA, *Escherichia coli* E. c., *Pseudomonas aeruginosa* P. a., *Staphylococcus epidermis* S. e., *Candida albicans* C. a., *Enterococcus hirae* E. h., *Aspergillus niger* A. n., *Aspergillus fumigatus* A. f., n. b. not defined.

TABLE 1









(part 1): MIC values (µg/ml)*						
Substance	Code	S.a.	MRSA	E.c.	P.a.	S.e.
	GG20-3	32	32	>64	>64	16

TABLE 1-continued

	GG20-4	4-8	8	>64	>64	4
	GG20-5	16	16	n. b.	n. b.	n. b.
	GG28	13 (3)	18 (3)	n. b.	n. b.	n. b.
Vancomycin			<2 (2)			
Tetracyclin			64			

(part 2): MIC values (µg/ml)*


Substance	Code	C. a. (Hefe)	E. h.	A. n.	A. f.
	GG20-3	64 10 (3)	n. b.	n. b.	n. b.
	GG20-4	4 11 (5)	16	64	8
	GG20-5	8	n. b.	n. b.	n. b.



[0016] Compounds with a chain length of $n=4$ were obtained analogously to the reaction with substituted 5-chloro-valerianophenon derivats with dimethylamine in the sense of a nucleophilic substitution.

[0017] The compound 2 was obtained based on the compound 1 in the sense of a Wolff-Kishner-Reduction following by the Huang-Minlon-Variant:

Scheme 2: Synthesis of 4-bromo-4'-(4-dimethylaminobutyl)biphenylhydrochloride 2 (GG28).



[0018] The basic biphenyls of the present invention represent high effective inhibitors of the growth of bacteria and fungi. They can be easily and economically synthesized by standard methods with high purity. Accordingly, the present invention provides compounds for inhibiting the growth of bacteria and/or fungi and, consequently, drugs for the treatment of infections of bacteria and/or fungi.

[0019] Studies of analogue compounds with a linker consisting of three, four and five carbon atoms also without ketofunction show that the antibacterial effect is dependent on the length of the side chain. Alternatively, the replacement of the N-dimethylamino group to a N-phenylpiperazino-group leads to loss of activity. In a test experiment a MIC

value for 4'-(4-bromophenyl)-4-(N-phenylpiperazino)butyrophenone hydrochloride could not be determined, since the substance did not inhibit the growth of *Staphylococcus aureus* also after addition of higher concentrations.

[0020] Pathogens in the growth phase are killed with lower concentrations as those which are in the dormant phase. In a test experiment, the test germs were once used in the exponential phase of their growth (18 h before incubation), compared to an inoculum which was harvested, and used after 48 h (quiescent period). Cells which were not as capable of dividing, showed a 2-3 fold higher MIC as the active cells. Thus, there are indications that quiescent phase cells are less sensitive.

[0021] As evident from the explanations and examples above, the substances of the present invention can also be used as laboratory reagents for inhibiting the growth of bacteria and/or fungi in cell culture, similar as described for the selection marker system neomycin and neomycin-phosphotransferase system.

[0022] The preparation in accordance with the present invention is illustrated by way of examples.

EXAMPLE 1

4'-(4-bromophenyl)-4-dimethylaminobutyrophenon-hydrochloride 1

[0023] 25.7 mmol (6 g) 4-bromobiphenyl, 25.7 mmol (3.6 g) 4-chloro butyryl chloride and 32.2 mmol (4.2 g) aluminumtrichloride are weighed out in a 250 mL piston before 50 mL dried dichloromethane are added. After stirring overnight at room temperature, the mixture is placed on ice and the precipitated aluminum hydroxide is brought into solution with concentrated hydrochloric acid. Subsequently, the organic phase is separated and extracted with water as long as the water phase shows a neutral pH value. The organic phase is then dried over Na_2SO_4 , and the solvent is filtered and removed under vacuum. The residuum is recrystallized from a mixture of cyclohexane/ethyl acetate.

[0024] 1.2 mmol (500 mg) NaI are added to 1.2 mmol (184 mg) of the resulting pure product in 5 mL ethylmethylketone and heated for 1 1/2 h at 90° C. in an oil bath. Subsequently, the solution is removed in a vacuum, 1 mL of dimethylamine as well as 30 mL absolute ethanol are added and heated for further 8 h at reflux.

[0025] After removing the solvent in the vacuum, ice is added to the residue and the precipitated raw product is aspirated. Recrystallization from cyclohexane/ethyl acetate results 1 in a yield of 82% (377 mg); mp=224° C.

[0026] $^1\text{H-NMR}$ Data (300 MHz, DMSO-d_6): 1.94 (m., 2H, $-\text{CH}_2-$), 2.8 (s, 6H, $\text{N}(\text{CH}_3)_2$), 3.15 (m, 4H, COCH_2- ; $-\text{CH}_2\text{NH}(\text{CH}_3)_2$), 7.70 (m, 4H, Ar-H); 7.85 (d, $^3\text{J}=8.7$ Hz, 2H, H-3', H-5'), 8.04 (d, $^3\text{J}=8.7$ Hz, 2H, H-2', H-6'), 9.21 (bs, 1H, NH). $^{13}\text{C-NMR}$ -Data (75 MHz, DMSO-d_6): 18.4 (C-3), 35.0 (C-2), 42.0 ($2\times\text{CH}_3$), 56.1 (C-4), 122.0 (C-4"), 126.8 (2C, C-3', C-5'), 128.6 (2C, C-2', C-6'), 129.0 (2C, C-2"; C-6"), 131.9 (2C, C-3", C-5"), 135.5 (C-1"), 138.0 (C-1'), 143.2 (C-4'), 198.2 (CO).

EXAMPLE 2

4-bromo-4'-(4-dimethylaminobutyl)biphenylhydrochloride 2

[0027] 0.02 mol (6.96 g), 1, 5 mL hydrazine monohydrate and 60 mL absolute ethanol are added in a 100 mL piston.

Subsequently, it is heated at the reflux until a complete solution occurs (12 h). Afterwards, 40 mL ethanol are removed under vacuum and 40 mL triglycol as well as 10 g of potassium hydroxide and 3 ml hydrazine monohydrate are added. The mixture is heated a further hour at 80° C. (development of gas!) before the temperature is up-regulated until the thermometer indicates 200 to 220° C. The mixture is cooled down and the resulting precipitation is aspirated and recrystallized from ethanol. Yield 85% (5.6 g); mp=245° C.

[0028] ¹H-NMR-Data (300 MHz, CDCl₃): 1.39 (m, 2H, —CH₂—), 1.62 (m, 2H, —CH₂—), 2.27 (s, 6H, N(CH₃)₂), 2.36 (t, ³J=8.4 Hz, 2H, —CH₂NH(CH₃)₂), 2.55 (t, ³J=8.4 Hz, 2H, Ar—CH₂—), 7.18 (d, ³J=8.7 Hz; 2H, H-2', H-6'); 7.42 (m, 4H, H-3', H-5', H-2'', H-6''), 7.48 (d, ³J=8.7 Hz; 2H, H-3'', H-5''). ¹³C-NMR-Data (75 MHz, CDCl₃): 27.4 (C-3), 29.0 (C-2), 35.7 (C-1), 45.9 (2×CH₃), 59.1 (C-4), 122.0 (C-4''), 127.7 (2C, C-3', C-5'), 128.7 (2C, C-2', C-6'), 130.1 (2C, C-2'', C-6''), 132.2 (2C, C-3'', C-5''), 133.7 (C-4'), 135.5 (C-1''), 137.6 (C-1').

[0029] The melting points of the synthesized substances were measured with the melting point apparatus Büchi 510 device and microhotplate Thermovar (Company Reichert). The NMR spectra were measured with the nuclearmagneticspectrometer Bruker ARX 300 and the IR spectra (as KBr-Presslinge) with a Perkin-Elmer FT-IR 16 PC spectrometer. The mass spectra were measured with a device of type Hewlett-Packard 5989. Elementary analysis was performed in the Institute of Inorganic Chemistry, of the CAU Kiel using a CHNS-analyzer of the company Hekatech GmbH. Unless otherwise stated, the chemicals including vancomycin-HCl and tetracycline-HCl at the highest purity were purchased at the company Sigma-Aldrich GmbH.

[0030] The minimum inhibitory concentrations of the substances of the present inventions against various infectious germs, i.a. against methicillin-resistant *Staphylococcus aureus* were determined with the Bouillon-microdilution-method in accordance with the procedure M07-A8 of the Clinical and Laboratory Standards Institute (Pennsylvania, USA).

Test Organisms

[0031] (Source: German Collection of Microorganisms and Cell Cultures, Braunschweig), *Enterococcus hirae* ATCC 10541, *Staphylococcus epidermidis* ATCC 12228, *Staphylococcus aureus* ATCC 6538, methicillin-resistant *Staphylococcus aureus* ATCC 33592, *Bacillus cereus* ATCC 11778, *Candida albicans* ATCC 10231, *Aspergillus niger* ATCC 16404, *Aspergillus fumigatus* ATCC 9197.

Test Vessels

[0032] Sterile 96-well microtiterplates of plastic with rounded bottoms of the test wells (wells; BRANDplates™, Ref. 781960).

Cultivation of Test Organisms

[0033] In sterile tubes with titled agar, bacteria with Tryptic Soy Agar (Merck, Art. No. 1.05458.05), incubation 18-24 h at 34° C., fungi with Sabouraud 4% Glucose Agar (Merck 1.05438.05), incubation *Candida albicans* 18-24 h at 34° C., mold fungous 7 days at 22° C.

[0034] The long-term cultures of the test organisms are incubated at titled agar at 22° C. and each inoculated after 4 weeks on fresh medium. After every fifth culture passage or

ascertainment of impurities to the correspondent culture is discarded and newly grown from the lyophilized.

Production of Inocula

[0035] Suspensions are produced on titled agar by floating with 0.9% sterile NaCl solution. The turbidity is adjusted photometrically by dilution according to the turbidity of the Mc Farland-standard 0.5. Subsequently, the bacterial suspensions are diluted in the ratio of 1:10, the Pi-suspension of the fungi remains undiluted.

[0036] The number of the germs in the inocula is adjusted so that there are approximately 5×10⁵ colony forming units per milliliter after inoculation of the test wells.

Preparing Stock Solutions of the Antibiotics

[0037] The amount of a substance to be tested in a trial is weighed out on a microbalance with an Eppendorf reaction vessel and dissolved in an appropriate volume of dimethyl sulfoxide so that the concentration of the resulting solution corresponds to the 21 times highest final test concentration. The stock solution is subsequently diluted in a serial in a ratio respectively 1 ad 2 with DMSO resulting in seven different concentrations in the wells of a predilutionplate. In the eighth well only DMSO is pipetted.

Loading the Plates

[0038] In each of the 96 wells of a test plate 95 µl sterile Mueller Hinton II medium (Cation adjusted, BBL™ Ref 212322) for bacteria and sterile Sabouraud 2% glucose medium (Difco™ Ref 238230) for fungus are added with an eight-channel pipette (Socorex 50-200 µL). 5 µL of the solution to be tested are pipetted in each of the test wells of the plate (8-channel pipette, Eppendorf-Research, 5-10 µL). They are transferred in this way from the pre-dilutions that the horizontal row A of the highest concentration and to row G which shows a half reduced concentration of the substance to be tested and in row H where only DMSO is pipetted. Subsequently, 5 µL of the inocula are pipetted in the test wells using an eight-channel pipette. In one test at least two, maximum four rows are inoculated with the same substance and the same germs. The concentrations of the test substance are each 128 µg/mL to 2 µg/mL.

Evaluation and Incubation of the Microtiter Plates

[0039] The prepared plates are converted to a microtiter plate reader (Anthos htIII) which is connected to a printer. In this device, the plates are shaken for 60 seconds with high frequency before each measurement and the absorption of the probes is measured at the wavelength of light of 590 nm.

[0040] After determination of the initial values the plates are incubated at 34° C. for 16 to 20 h for bacteria and *Candida albicans*, and 68 to 72 h at 34° C. for mould fungus.


[0041] Following this the microtiterplates are measured again as described above.

Determination of the Minimum Inhibitory Concentration (MIC)

[0042] The minimum inhibitory concentration is the lowest concentration of the tested substance which is able to inhibit the complete growth of the particular test organism, i.e. where after incubation no turbidity is measurable. The particular

determination is valid whether in the growth control (DMSO and medium) a distinct turbidity is observed.

1. A substance of the general formula



for use as antibacterial and/or antifungal drug, wherein

X is a methylene or a carbonyl group;

R₁, R₂ and R₃ are each selected from the group consisting of hydrogen, an alkyl group having a chain length of 1-4 carbon atoms, an alkoxy group with a chain length of 1-3 carbon atoms, and a halogen;

R₄ and R₅ are each selected from the group consisting of hydrogen and an alkyl group with a chain length of 1-4 carbon atoms; and

n=3 to 6.

2. The substance of claim 1, wherein the substance is available as a hydrochloride, a salt, or as a tautomer.

3. The substance of claim 1, wherein the alkyl group is a methyl (CH₃)—, an ethyl (CH₂CH₃)—, a propyl (CH₂CH₂CH₃)—, an isopropyl (CH(CH₃)CH₂)—, a butyl (CH₂CH₂CH₂CH₃)—, an isobutyl (CH(CH₃)CH₂CH₂)— or a tertiary butyl (C(CH₃)₃)— group.

4. The substance of claim 1, wherein the alkoxy group is a methoxy (OCH₃)—, an ethoxy (OCH₂CH₃)—, a propoxy (OCH₂CH₂CH₃)— or an isopropoxy (OCH(CH₃)CH₂)— group.


5. The substance of claim 1, wherein the halogen is fluorine (F), chlorine (Cl), bromine (Br), or iodine (I).

6. The substance of claim 1, wherein the water-solubility of the substance is equal to or greater than 1,000 mg/l.


7. The substance of claim 1, wherein the water-solubility of the substance is equal to or greater than 5,000 mg/l.

8. The substance of claim 1, wherein the water-solubility of the substance is equal to or greater than 10,000 mg/l.

9. The substance of claim 1, wherein the substance is 4'(4-bromophenyl)-4-dimethylaminobutyrophenonylhydrochloride with the structural formula



10. The substance of claim 1, wherein the substance is 4-brom-4'-(4-dimethylaminobutyl)biphenylhydrochloride with the structural formula



* * * * *